

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

BI

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K	A2	(11) International Publication Number: WO 00/01347 (43) International Publication Date: 13 January 2000 (13.01.00)
(21) International Application Number: PCT/US99/13925 (22) International Filing Date: 21 June 1999 (21.06.99) (30) Priority Data: 60/091,658 2 July 1998 (02.07.98) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): SNYDER, Daniel, Earl [US/US]; 7348 Yorkshire Boulevard, Indianapolis, IN 46229 (US). (74) Agents: GRAY, Edouard, P. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AE, AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: FORMULATIONS FOR CONTROLLING HUMAN LICE (57) Abstract Safer pediculicidal formulations comprising a spinosyn, or a physiologically acceptable derivative or salt thereof, and a physiologically acceptable carrier, and methods of controlling lice infestations in a human with these formulations are provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

FORMULATIONS FOR CONTROLLING HUMAN LICE

5 Infestation of the human body by lice is an
increasingly prevalent social and health problem in many
countries, including the United States. Lice are very small
insects (about 2-3mm in length). They deposit eggs either
on a hair or fabric fiber and attach them firmly with a
cementlike excretion. The eggs generally hatch in about six
10 to ten days, depending on temperature. The empty shells
remaining after the nymphs emerge from the eggs look like
white grains of sand. These shells are called nits.

15 The Anoplura, or sucking lice, are parasites found
on nearly all groups of mammals. Of the 15 recognized
families of Anoplura, two families, *Pediculidae* and
Pthiridae, have species found on humans. *Pediculus humanus*
is the only species in the family *Pediculidae* that infests
humans. It includes the head louse, *Pediculus humanus*
20 *capitis*; and the body or clothing louse, *Pediculus humanus*
humanus, sometimes called *Pediculus corporis*. The crab
louse, *Pthirus pubis*, is a distinct species and is the only
member of the family *Pthiridae* that infests humans. As used
herein, the term "human lice or louse" includes a member of
25 *Pediculus humanus* or *Pthirus pubis*.

-2-

Human lice are spread by crowding and common usage of clothing and combs. Initially, infestations result at most in irritation, but the irritation can lead to infection of the irritated area. There are at least three major diseases that are primarily transmitted by lice: epidemic typhus, trench fever and relapsing fever.

Although the human lice varieties are related, each of them has specific characteristics with regard to habitat and feeding. For example, head lice are small hard-shelled ectoparasites which cling to hair shafts while feeding, mating and laying eggs. The louse must remain on the head or it will die within a short period of time. Head lice proliferate at an incredible rate. A louse is ready to mate and reproduce within 10 hours after hatching. Under ideal conditions, a female louse may produce up to 300 eggs in its lifetime. Ideal conditions include an adequate food supply, environmental temperatures from about 28°C. to about 32° C., and relative humidity of about 70% to about 90%.

Poor hygienic and grooming habits are also known to contribute significantly to the spread of lice. Thus, lice infestations are most serious in geographical areas where the inhabitants have both substandard hygienic

-3-

facilities and practices. Lice can be a problem, however, even when conditions are relatively sanitary.

5 The louse's hard chitinous exoskeleton serves
as protection from external elements. Lice eggs (or ova)
are similarly protected by a chitinous sheath surrounding
the eggs and attached to the hair shaft. Although lice may
be affected by the use of an insecticide, the eggs often
remain resistant to attack. Thus, optimum treatment of a
10 lice infestation includes both a pediculicide, which kills
the adult lice, and an ovicide, which interrupts the
development of the eggs.

15 Biologically active agents have been used for
some time in attempts to control lice. For example, lindane
(gamma-benzene hexachloride), organophosphates (malathion),
natural pyrethrins, and synthetic compounds known as
pyrethroids (such as permethrin) have been used as
pediculicides in lice treatment formulations. These agents
20 however, have drawbacks. For example, lindane has a poor
safety profile, and lice have developed resistance to it.
Natural pyrethrin requires frequent follow-up treatments
because it provides only short term residual action.
Synthetic pyrethroids, although more effective against lice
25 than natural pediculicides, are often more toxic to the
subject being treated.

-4-

Spinosyns (also known as A83453 factors) are agricultural insecticides that have shown activity against 1) southern armyworm and other insects in the order *Lepidoptera*, 2) cotton aphid and other members of the order *Homoptera*, and 3) stable flies, blow flies and mosquitos, which are members of the insect order *Diptera*. (See U.S. Patent No. 5,362,634, *infra*). Spinosyn A has an excellent human and animal safety and toxicological profile.

This invention relates to formulations for controlling a lice infestation in a human comprising a spinosyn, or a physiologically acceptable derivative or salt thereof, and a physiologically acceptable carrier. The invention further relates to methods of controlling a lice infestation in a human comprising topically administering to the human an amount of a spinosyn, or a physiologically acceptable derivative/salt thereof, that controls the lice. The formulations and methods of this invention are safer and more effective than those presently available. A particular benefit of the formulations is their effectiveness against louse species that have become resistant to currently used products. Preferred formulations and methods of this invention are hair care formulations, such as shampoos, lotions and conditioners, and methods of using these hair care formulations for controlling a lice infestation in a human.

-5-

Especially useful pediculicidal/ovicidal (anti-lice) formulations of this invention are hair-care formulations. Especially useful hair-care formulations are shampoos.

5 In another aspect this invention provides the use of a spinosyn, or a physiologically acceptable derivative or salt thereof, or a formulation containing either an spinosyn or derivative or salt thereof, for the manufacture of a medicament for controlling lice in a human.

10 The term "controlling a lice infestation" refers to treating an active lice infestation or preventing an infestation in a human who is likely to be exposed to a lice infestation.

15 Spinosyns are naturally derived fermentation products. They are macrolides produced by cultivation of *Saccharopolyspora spinosa*. The fermentation produces multifactors, including spinosyn A and spinosyn D (also
20 called A83543A and A83543D). Spinosyn A and spinosyn D are the two spinosyns that are most active as insecticides. A product comprised mainly of these two spinosyns (approximately 85% A and 15% D) is available commercially from Dow Agrosiences under the name spinosad. The name
25 "spinosad" comes from a contraction of the spinosyns "A" and "D".

-6-

Each spinosyn has a 12-membered macrocyclic ring that is part of an unusual tetracyclic ring system to which two different sugars are attached, the amino-sugar forosamine and the neutral sugar 2N,3N,4N-tri-O-methylrhamnose. This unique structure sets the spinosyns apart from other macrocyclic compounds.

Spinosyn A (A83543A) was the first spinosyn isolated and identified from the fermentation broth of *Saccharopolyspora spinosa*. Subsequent examination of the fermentation broth revealed that the parent strain of *S. spinosa* produced a number of spinosyns that have been labeled A to J (A83543A to J). Compared to spinosyn A, spinosyns B-J are characterized by differences in the substitution patterns on the amino group of the forosamine, at selected sites on the tetracyclic ring system and on 2N,3N,4N-tri-O-methylrhamnose. The strains of *S. spinosa* currently in use produce a mixture of spinosyns of which the primary components are spinosyn A (~85%) and spinosyn D (~15%). Additional spinosyns, lettered from K to W, have been identified from mutant strains of *S. spinosa*.

The term "spinosyn or a derivative thereof" as used herein refers to an individual spinosyn factor (A, B, C, D, E, F, G, H, J, K, L, M, N, O, P, Q, R, S, T, U, V, W or Y), an N-demethyl derivative of an individual spinosyn

-7-

factor, or a combination thereof. For convenience, the term "spinosyn component" will also be used herein to mean an individual spinosyn, or a physiologically acceptable derivative or salt thereof, or a combination thereof.

5

10

15

20

25

Boeck et al. described spinosyns A-H and J (which they called A83543 factors A, B, C, D, E, F, G, H and J), and salts thereof, in U.S. Patent Nos. 5,362,634 (issued Nov. 8, 1994); 5,496,932 (issued March 5, 1996); and 5,571,901 (issued Nov. 5, 1996). Mynderse et al. described spinosyns L-N (which they called A83543 factors L, M and N), their N-demethyl derivatives, and salts thereof, in U.S. Patent No. 5,202,242 (issued Apr. 13, 1993); and Turner et al. described spinosyns Q-T (which they called A83543 factors Q, R, S and T), their N-demethyl derivatives, and salts thereof, in U.S. Patent Nos. 5,591,606 (issued January 7, 1997) and 5,631,155 (issued May 29, 1997). These patents are incorporated herein by reference. Spinosyns K, O, P, U, V, W and Y are described, for example, by Carl V. DeAmicis, James E. Dripps, Chris J. Hatton and Laura I. Karr in American Chemical Society's Symposium Series: Phytochemicals for Pest Control, Chapter 11, "Physical and Biological Properties of Spinosyns: Novel Macrolide Pest-Control Agents from Fermentation", pages 146-154 (1997).

-8-

The spinosyns can react to form salts. Salts that are physiologically acceptable are also useful in the formulations and methods of this invention. The salts are prepared using standard procedures for salt preparation.

5 For example, spinosyn A can be neutralized with an appropriate acid to form an acid additional salt. The acid addition salts of spinosyns are particularly useful.

Representative suitable acid addition salts include salts formed by reaction with either an organic or inorganic acid
10 such as, for example, sulfuric, hydrochloric, phosphoric, acetic, succinic, citric, lactic, maleic, fumaric, cholic, pantoic, mucic, glutamic, camphoric, glutaric, glycolic, phthalic, tartaric, formic, lauric, stearic, salicylic, methanesulfonic, benzenesulfonic, sorbic, picric, benzoic,
15 cinnamic and like acids.

In addition to the spinosyn component, the formulations of this invention may further include one or more other compounds that have activity against lice such
20 as, for example, synthetic pyrethroids, natural pyrethrins, and lindane. All ratios, percentages, and parts discussed herein are "by weight" unless otherwise specified.

The anti-lice formulations of this invention
25 may be formulated in a number of ways. Particularly useful formulations are shampoos, conditioners, and lotions. These

-9-

formulations optionally also comprise one or more of the following: a) a surfactant; b) from about 1% to about 10% of a non-volatile silicone material; and/or c) from about 0.5% to about 5% of a suspending agent.

5

The shampoo formulations of this invention comprise a spinosyn, or a physiologically acceptable derivative or salt thereof, together with water, a surfactant, and an amide and may optionally comprise another anti-lice agent, a silicone compound, a suspending agent and other cosmetically acceptable components.

10

Human hair becomes soiled due to contact with the surrounding atmosphere and the build up of sebum secreted by the head. When the hair is soiled, it has a dirty feel and an unattractive appearance. The shampoo formulations of this invention both clean the hair and effectively control lice infestation.

15

When used in a shampoo formulation, the spinosyn component is present at a level of from about 0.1% to about 30%, preferably from about 1% to about 10%.

20

Surfactants suitable for use in these formulations can be any of a wide variety of synthetic anionic, amphoteric, zwitterionic and non-ionic surfactants.

25


-10-

Surfactants are generally present in shampoo formulations at a level of from about 5% to about 30%, preferably from about 15% to about 25%.

5 Examples of synthetic anionic surfactants are the alkali metal salts of organic sulfuric reaction products having an alkyl radical containing from 8-22 carbon atoms and a sulfonic acid or sulfuric acid ester radical (included in the term alkyl is the alkyl portion of higher acyl
10 radicals). Sodium, ammonium, potassium or triethanolamine alkyl sulfates are preferred, especially those obtained by sulfating the higher alcohols (C_8-C_{18} carbon atoms); sodium coconut oil fatty acid monoglyceride sulfates and sulfonates; sodium or potassium salts of sulfuric acid
15 esters of the reaction product of 1 mole of a higher fatty alcohol (e.g., tallow or coconut oil alcohols) and 1 to 12 moles of ethylene oxide; sodium or potassium salts of alkyl phenol ethylene oxide ether sulfate with 1 to 10 units of ethylene oxide per molecule and in which the alkyl radicals
20 contain from 8 to 12 carbon atoms; sodium alkyl glyceryl ether sulfonates; the reaction product of fatty acids having from 10 to 22 carbon atoms esterified with isethionic acid and neutralized with sodium hydroxide; and water soluble salts of condensation products of fatty acids with
25 sarcosine.

-11-

Examples of zwitterionic surfactants are derivatives of aliphatic quaternary ammonium, phosphonium, and sulfonium compounds, in which the aliphatic radicals can be straight or branched, and wherein one of the aliphatic substituents contains from about 8 to 18 carbon atoms and one contains an anionic water-solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. A general formula for these compounds is:



wherein R^2 contains an alkyl, alkenyl, or hydroxyalkyl radical of from about 8 to about 18 carbon atoms, from 0 to about 10 ethylene oxide moieties and from 0 to 1 glyceryl moiety; Y is a nitrogen, phosphorus, or sulfur atom; R^3 is an alkyl or monohydroxyalkyl group containing 1 to about 3 carbon atoms; x is 1 when Y is sulfur and 2 when Y is nitrogen or phosphorus; R^4 is alkylene or hydroxyalkylene of from 1 to about 4 carbon atoms, and Z is a carboxylate, sulfonate, sulfate, phosphonate, or phosphate radical.

Examples include:

4-[N,N-di(2-hydroxyethyl)-N-octadecylammonio]-butane-1-carboxylate;

-12-

5-[S-3-hydroxypropyl-S-hexadecylsulfonio]-3-hydroxy-pentane-1-sulfate;

3-[P,P-diethyl-P-3,6,9-trioxatetradecylphosphonio]-2-hydroxypropane-1-phosphate;

5 3-[N,N-dipropyl-N-3-dodecoxy-2-hydroxypropylammonio]-propane-1-phosphate;

3-(N,N-dimethyl-N-hexadecylammonio)propane-1-sulfonate;

10 3-(N,N-dimethyl-N-hexadecylammonio)-2-hydroxypropane-1-sulfonate;

N,N-di(2-hydroxyethyl)-N-(2-hydroxydodecyl)ammonio]-butane-1-carboxylate;

3-[S-ethyl-S-(3-dodecoxy-2-hydroxypropyl)sulfonio]-propane-1-phosphate;

15 3-[P,P-dimethyl-P-dodecylphosphonio]-propane-1-phosphonate; and

5-(N,N-di(3-hydroxypropyl)-N-hexadecylammonio)-2-hydroxypentane-1-sulfate.

20 Other zwitterionic surfactants, such as betaines, are also useful in the formulations of this invention. Examples of betaines include the higher alkyl betaines, such as coco dimethyl carboxymethyl betaine, lauryl dimethyl carboxymethyl betaine, lauryl dimethyl
25 alpha-carboxyethyl betaine, cetyl dimethyl carboxymethyl betaine, lauryl bis-(2-hydroxyethyl) carboxymethyl betaine,

-13-

stearyl bis-(2-hydroxypropyl) carboxymethyl betaine, oleyl dimethyl gamma-carboxypropyl betaine and lauryl bis-(2-hydroxypropyl) alpha-carboxyethyl betaine. The sulfobetaines may be represented by coco dimethyl

5 sulfopropylbetaine, stearyl dimethyl sulfopropyl betaine, lauryl dimethyl sulfoethyl betaine, lauryl bis-(2-hydroxyethyl) sulfopropyl betaine, and the like. Amido betaines and amidosulfobetaines, wherein an $\text{RCONH}(\text{CH}_2)_3$ radical is attached to the nitrogen atom of the betaine, are

10 also useful in the formulations of this invention.

Examples of amphoteric surfactants that can be used in the formulations of this invention are those which are derivatives of aliphatic secondary or tertiary

15 amines in which the aliphatic radical is straight or branched and wherein one of the aliphatic substituents contains from about 8 to about 18 carbon atoms and one contains an anionic water solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. Examples of

20 amphoteric surfactants are sodium 3-dodecylaminopropionate, sodium 3-dodecylaminopropane sulfonate, N-alkyltaurines such as the one prepared by reacting N-dodecylamine with sodium isethionate (see U.S. Pat. 2,658,072, Example 3), N-higher alkyl aspartic acids (see U.S. Pat. 2,438,091), and products

25 sold under the trade name "Miranol" and described in U.S. Pat. 2,528,378.

-14-

Nonionic surfactants, which are preferably used in combination with an anionic, amphoteric or zwitterionic surfactant, are compounds produced by the condensation of alkylene oxide groups (hydrophilic in nature) with an organic hydrophobic compound, which may be aliphatic or alkyl aromatic in nature. Examples of nonionic surfactants include:

1) Polyethylene oxide condensates of alkyl phenols, e.g., the condensation products of alkyl phenols having an alkyl group containing from about 6 to 12 carbon atoms in either a straight or branched chain configuration, with ethylene oxide, the ethylene oxide being present in amounts equal to 10 to 60 moles of ethylene oxide per mole of alkyl phenol. The alkyl substituent in these compounds may be derived from polymerized propylene, diisobutylene, octane, or nonane, for example.


2) Condensates of ethylene oxide with a product of the reaction of propylene oxide and ethylene diamine products which may be varied in formulation depending upon the desired balance between the hydrophobic and hydrophilic elements. For example, compounds containing from about 40% to about 80% polyoxyethylene by weight and having a molecular weight of from about 5,000 to about 11,000 resulting from the reaction of ethylene oxide groups with a hydrophobic base comprising the reaction product of

-15-

ethylene diamine and excess propylene oxide, and having a molecular weight on the order of 2,500 to 3,000, are satisfactory.

5 3) The condensation product of aliphatic alcohols having from 8 to 18 carbon atoms, in either straight or branched chain configuration, with ethylene oxide, e.g., a coconut alcohol ethylene oxide condensate having from 10 to 30 moles of ethylene oxide per mole of
10 coconut alcohol, the coconut alcohol fraction having from 10 to 14 carbon atoms.

 4) Long chain tertiary amine oxides corresponding to the following general formula:




wherein R_1 contains an alkyl, alkenyl, or monohydroxy alkyl radical of from about 8 to about 18 carbon atoms, from 0 to
20 about 10 ethylene oxide moieties, and from 0 to 1 glyceryl moiety, and R_2 and R_3 contain from 1 to about 3 carbon atoms and from 0 to about 1 hydroxy group, e.g., methyl, ethyl, propyl, hydroxyethyl, or hydroxypropyl radicals. The arrow in the formula represents a semipolar bond. Examples of
25 amine oxides suitable for use in these formulations include dimethyldodecylamine oxide, oleyldi(2-hydroxyethyl)amine

-16-

oxide, dimethyloctylamine oxide, dimethyl-decylamine oxide,
dimethyl-tetradecylamine oxide, 3,6,9-
trioxaheptadecyldiethylamine oxide, di(2-hydroxyethyl)-
tetradecylamine oxide, 2-dodecoxyethyldimethylamine oxide,
5 3-dodecoxy-2-hydroxypropyldi(3-hydroxypropyl) amine oxide,
and dimethylhexadecylamine oxide.

5) Long chain tertiary phosphine oxides of
the following general formula:



wherein R contains an alkyl, alkenyl or monohydroxy-alkyl
radical of from about 8 to 18 carbon atoms, from 0 to about
10 ethylene oxide moieties and from 0 to 1 glyceryl moiety,
15 and R' and R'' are each alkyl or monohydroxyalkyl groups
containing from 1 to 3 carbon atoms. The arrow in the
formula represents a semipolar bond.

20 Examples of suitable phosphine oxides
include: dodecyldimethyl-phosphine oxide,
tetradecyldimethylphosphine oxide,
tetradecylmethylethylphosphine oxide, 3,6,9-
trioxaoctadecyldimethylphosphine oxide,
25 cetyldimethylphosphine oxide, 3-dodecoxy-2-
hydroxypropyldi(2-hydroxyethyl) phosphine oxide,

-17-

stearyldimethylphosphine oxide, cetylethylpropylphosphine oxide, oleyldiethylphosphine oxide, dodecyldiethylphosphine oxide, tetradecyldiethylphosphine oxide, dodecyldipropylphosphine oxide, dodecyldi(hydroxymethyl)phosphine oxide, dodecyldi(2-hydroxyethyl)phosphine oxide, tetradecylmethyl-2-hydroxypropylphosphine oxide, oleyldimethylphosphine oxide, 2-hydroxydodecyldimethylphosphine oxide.

6) Long chain dialkyl sulfoxides containing one short chain alkyl or hydroxyalkyl radical of 1 to 3 carbon atoms (usually methyl) and one long hydrophobic chain which contains alkyl, alkenyl, hydroxyalkyl, or keto alkyl radicals containing from about 8 to about 20 carbon atoms, from 0 to about 10 ethylene oxide moieties and from 0 to 1 glyceryl moiety. Examples include: octadecyl methyl sulfoxide, 2-ketotridecyl methyl sulfoxide, 3,6,9-trioxaoctadecyl 2-hydroxyethyl sulfoxide, dodecyl methyl sulfoxide, oleyl 3-hydroxypropyl sulfoxide, tetradecyl methyl sulfoxide, 3-methoxytridecyl methyl sulfoxide, 3-hydroxytridecyl methyl sulfoxide, and 3-hydroxy-4-dodecoxybutyl methyl sulfoxide.

Many additional nonsoap surfactants are described in *McCutcheon's Detergents and Emulsifiers*, 1998 Annual, published by M.C. Publishing Company, Inc.;

-18-

McCutcheon Division, 175 Rock Rd., Glen Rock, NJ, 07425,
U.S.A.

5 Anionic surfactants, particularly the alkyl
sulfates, the ethoxylated alkyl sulfates and mixtures
thereof, as well as the amido betaines, are preferred for
use in the shampoo formulations of this invention.

10 Amides enhance the lathering of the
formulations by emulsifying the shampoo components and the
active component(s). The amides used in the present
formulations can be any of the alkanolamides of fatty acids
known for use in shampoos. These are generally mono- and
diethanolamides of fatty acids having from about 8 to about
15 14 carbon atoms. Other suitable amides are those having
multiple ethoxy groups such as PEG-3 lauramide.

20 In the shampoo formulations, the amide is
generally present at a level of about 1% to about 7%,
preferably from about 2% to about 5%, of the formulation.
Preferred amides are coconut monoethanolamide, coconut
diethanolamide, and mixtures thereof.

25 The shampoo formulations of this
invention also contain water. Water is typically present in
the shampoos at levels of from about 50% to about 80%,

-19-

preferably from about 60% to about 75%. After adding water, the relative viscosity of the formulation is generally in the range of from about 4,000 centipoise (cp) to about 25,000 cp, preferably from about 4,000 cp to about 12,000 cp, most preferably from about 4,000 cp to about 5,500 cp, measured at 1 RPM at 26.7° for 3 minutes using a Wells-Brookfield viscometer Model DV-CP-2 DVII, Model Cone CP-41. Viscosity modifiers and hydrotropes may be included to bring the formulation's viscosity within these ranges.

The shampoo formulations may also include one or more optional components such as silicone compounds; suspending agents and components that make the formulation more cosmetically acceptable.

Silicone compounds condition the hair and facilitate removal of the dead lice, their eggs and nits. Non-volatile silicone materials are used at levels from about 1% to about 10% of the formulations. Examples of useful silicone compounds are disclosed in U.S. Patent No. 5,292,504, Cardin et al., issued March 8, 1994, incorporated herein by reference.

Non-volatile silicone-containing compounds are preferred and are used at levels of from about 0.1% to about 10%, preferably from about 0.25% to about 3%, by weight of the formulation. Examples of non-volatile

-20-

silicones are polyalkyl siloxanes, poly alkylaryl siloxanes, polyether siloxane copolymers and mixtures thereof.

5 Useful polyalkyl siloxanes include, for example, polydimethyl siloxanes (PDMS) with viscosities ranging from about 5 to 15,000,000 cp at 25°. These siloxanes are available, for example, from the General Electric Company as the Viscasil series and from Dow Corning as the Dow Corning 200 series. The viscosity can be
10 measured by means of a glass capillary viscometer as set forth in Dow Corning Corporate Test Method CTM0004, Jul. 20, 1970.

15 Useful polyalkylaryl siloxanes include polymethylphenyl siloxanes having viscosities of from about 5 to about 15,000,000 cp at 25°. These siloxanes are available, for example, from the General Electric Company as SF 1075 methyl phenyl fluid or from Dow Corning as 556 Cosmetic Grade Fluid.

20 Useful polyether siloxane copolymers include polypropylene oxide modified polydimethylsiloxanes (available, for example, from Dow Corning as DC-1248), ethylene oxide or mixtures of ethylene oxide and propylene
25 oxide. Water insoluble ones are most useful.

-21-

The siloxanes are able to condition the hair due to their ability to lubricate the hair, providing wet and dry combing benefits. Viscous, higher molecular weight siloxanes provide the best conditioning benefits and are, therefore, preferred. Fluids and gums of the siloxane polymers are most desirable. Siloxane polymer gums are rigid as opposed to a liquid or fluid, with high mass molecular weights of from about 200,000 to about 1,000,000 as viscosities from about 100,000 cp to about 150,000,000 cp at 25°C. Such gums are discussed in U.S. Patent No. 5,292,504 (supra).

Suspending agents may be included to improve long term stability. Useful suspending agents include fatty amphiphilic crystalline materials having needle-like or platelet structures, polymeric materials, clays, fumed metal oxides, and mixtures thereof. These agents are known in the art (see U.S. Patent No. 5,292,504).

Suitable crystalline amphiphilic materials are those that have needle or platelet-type structures. Such compounds include long chain (C_{16} - C_{22}) acyl derivatives, such as ethylene glycol esters of fatty acids (e.g., ethylene glycol distearate); long chain (C_{16} - C_{22}) alkanol amides of fatty acids, such as stearamide MEA, stearyl stearate, and distearyl dithiopropionate; and mixtures thereof.

-22-

Polymeric materials that are useful as suspending agents include cross-linked polyacyclic acids (such as the Carbopol series, available from the B.F. Goodrich Chemical Company), guar gum and its derivatives, xanthan gum, cross linked copolymers of ethylene/maleic anhydrides, and mixtures thereof.

Clays and fumed metal oxides are also effective suspending agents. Examples include magnesium aluminum silicates (such as the Veegum series, available from R. T. Vanderbilt Company, Inc.), sodium aluminum silicates (such as the Laponite series, available from Laponite United States), fumed silica, fumed alumina, fumed titania and mixtures thereof.

In the shampoo formulations of this invention suspending agents are generally present in amounts of from about 0.5% to about 5%, preferably from 0.5% to about 3%. The long chain acyl derivatives such as ethylene glycol esters of fatty acids are preferred. Most preferred is ethylene glycol distearate.

Other optional components that can improve the cosmetic acceptability of the formulations acceptable are known in the art and include, e.g., preservatives, such as methyl paraben, propyl paraben, methylisothiazolinone and imidazolidinyl urea; thickeners and viscosity modifiers,

-23-

such as amine oxides, block polymers of ethylene oxide and propylene oxide (such as Pluronic F88 offered by BASF Wyandotte), fatty alcohols (such as cetearyl alcohol), sodium chloride, ammonium chloride, sodium sulfate, polyvinyl alcohol, propylene glycol, and ethyl alcohol; hydrotropes, such as xylene sulfonate; pH adjusting agents, such as citric acid, succinic acid, phosphoric acid, sodium hydroxide, and sodium carbonate; perfumes, dyes, quaternary ammonium compounds, such as Polyquaternium 41, sequestering agents, such as disodium ethylenediamine tetraacetate; and pearlescing agents, such as distearic acid ester of ethylene glycol, stearic acid and palmitic acid diesters of polyethylene glycol, and stearic acid monoethanolamine. Generally, these optional components are used individually at a level of from about 0.1% to about 10% of the formulation.

The shampoo formulations of this invention are used in a conventional manner for cleaning hair. From about 10 g to about 30 g of a formulation is applied to wet hair and worked through both hair and scalp. The formulation is left on the hair and scalp for approximately 6-10 minutes and then is removed by rinsing. This process is repeated until the hair is clean.

-24-

A useful pediculicidal shampoo of this invention comprises:

- (a) from about 0.1% to about 10% of a spinosyn, or a physiologically acceptable derivative or salt thereof;
- (b) from about 5% to about 30% of a synthetic surfactant;
- (c) from about 1% to about 7% of an amide;
- and
- (d) water.

The hair conditioning formulations of this invention comprise a spinosyn component and a conditioner and may optionally comprise another anti-lice agent, such as permethrin or lindane. These conditioner formulations may also be used effectively to treat a lice infestation.

Hair conditioners are products that improve the appearance, feel and manageability of hair. Conditioners are particularly important when the hair has been damaged by treatments such as permanent waving, dyeing, teasing, and bleaching, or by atmospheric conditions, such as sunlight, that cause photo-catalyzed oxidation. These factors may cause hair to have poor texture, making it difficult to manage and comb, whether wet or dry.

-25-

When used in a hair conditioner formulation, the spinosyn component is present at a level of from about 0.1% to about 30%, preferably from about 1% to about 10%.

5 Conditioning products are well known and include "rinse-type" products, which are rinsed off shortly after being applied to clean hair, and "deep conditioners" which remain on the hair for extended periods of time.

10 One group of conditioners useful in the hair conditioner formulations of this invention are long chain quaternary ammonium compounds combined with lipid materials, such as fatty alcohols (see U.S. Pat. No. 3,155,591, Hilfer, issued Nov. 3, 1964, and U.S. Pat. No. 4,269,824,
15 Villamarin, et al., issued May 26, 1981). Another group of conditioners are lipids and quaternary ammonium compounds. These conditioners are used to form gel-type conditioner products having good in-use cosmetic and rheological characteristics. These types of gel-type formulations are
20 generally described in the following documents: Barry, "The Self Bodying Action of the Mixed Emulsifer Sodium Docecyl Sulfate/Cetyl Alcohol", *J. of Colloid and Interface Science*, **28**, 82-91 (1968); Barry et al., "The Self-Bodying Action of Alkyltrimethylammonium Bromides/Cetostearyl Alcohol Mixed
25 Emulsifiers; Influence of Quarternary Chain Length", *J. of Colloid and Interface Science*, **35**, 689-708 (1971); and Barry

-26-

et al., "Rheology of Systems Containing Cetomacrogol 1000 - (cetostearyl alcohol), I. Self-Bodying Action", *J. of Colloid and Interface Science*, **38**, 616-625 (1972).

5 The lipid materials used in these conditioners are present at a level of from about 0.5% to about 3%. These lipids are essentially water-insoluble, and contain hydrophobic and hydrophilic moieties. They include natural and synthetically-derived fatty materials selected from acids, acid derivatives, alcohols, esters, ethers, ketones, amides, and mixtures thereof, having alkyl chain lengths
10 from about 12 to about 22 carbon atoms, preferably from 16 to 18 carbon atoms in length. Fatty alcohols and fatty esters are preferred.

15 Useful fatty alcohols are known (see, for example, U.S. Pat. No. 3,155,591, supra; U.S. Pat. No. 4,165,369 (Watanabe et al., issued May 26, 1981); British Patent Specification 1,532,585, published Nov. 15, 1978; Fukushima et al., "The Effect of Cetostearyl Alcohol in Cosmetic Emulsions",
20 *Cosmetics & Toiletries*, **98**, 89-102 (1983); and Hunting, *Encyclopedia of Conditioning Rinse Ingredients*, at 204 (1987). Fatty alcohols are C₁₂-C₁₆ alcohols selected from cetearyl alcohol, cetyl alcohol, isostearyl alcohol, lanolin alcohol, lauryl alcohol, oleyl alcohol, stearyl alcohol, and
25 mixtures thereof. Preferred are cetyl alcohol, stearyl alcohol, and mixtures thereof. A particularly preferred

-27-

fatty alcohol is comprised of a mixture of cetyl alcohol and stearyl alcohol containing from about 55% to about 65% (by weight of mixture) of cetyl alcohol.


5 Useful fatty esters are also known (see Kaufman, et al., U.S. Pat. No. 3,341,465, issued Sep. 12, 1967). Fatty esters are fatty acids in which the active hydrogen has been replaced by the alkyl group of a monohydric alcohol. The monohydric alcohols are fatty alcohols as
10 described, supra. The fatty esters useful in these conditioner formulations include cetyl lactate, cetyl octanoate, cetyl palmitate, cetyl stearate, glyceryl monostearate, glyceryl laurate, glyceryl myristate, glyceryl oleate, glyceryl stearate, glyceryl monoacetate, and
15 mixtures thereof. Cetyl palmitate and glycerol monostearate, or mixtures thereof, are preferred.

Cationic surfactants may be used in these conditioning formulations, either singly or in combination,
20 generally at a level of from about 0.1% to about 5% of the final formulation. These surfactants contain amino or quaternary ammonium hydrophilic moieties that are positively charged when dissolved in the aqueous formulations of this invention. These cationic surfactants are known in the art
25 (see *McCutcheon's Detergents & Emulsifiers*, supra; Schwartz et al., *Surface Active Agents, Their Chemistry and*

-28-

Technology, New York: Interscience Publishers, 1949; U.S. Pat. No. 3,155,591, supra; U.S. Pat. No. 3,929,678 (Laughlin et al., issued Dec. 30, 1975); U.S. Pat. No. 3,959,461 (Bailey et al., issued May 25, 1976); and U.S. Pat. No. 4,387,090 (Bolich, Jr., issued Jun. 7, 1983).


Useful quaternary ammonium cationic surfactant materials are those of the general formula:



wherein R_1 is hydrogen, an aliphatic group of from 1 to 22 carbon atoms, or an aromatic, aryl or alkylaryl group having from 12 to 22 carbon atoms; R_2 is an aliphatic group having from 1 to 22 atoms; R_3 and R_4 are each alkyl groups having from 1 to 3 carbon atoms, and X is an anion selected from halogen, acetate, phosphate, nitrate and alkyl sulfate radicals. The aliphatic groups may contain ether linkages, and other groups such as amido groups, in addition to carbon and hydrogen atoms.

Other useful quaternary ammonium salts have the formula:

-29-



wherein at least one, but no more than 3, of the R groups is
 an aliphatic group having from 16 to 22 carbon atoms, and
 the remaining R groups are selected from hydrogen and alkyl
 groups having from 1 to 4 carbon atoms, and X is an ion
 selected from halogen, acetate, phosphate, nitrate and alkyl
 sulfate radicals. Tallow propane diammonium dichloride is an
 example of this type of quaternary ammonium salt.

Quaternary ammonium salts useful herein also
 include dialkyldimethylammonium chlorides wherein the alkyl
 groups have from 12 to 22 carbon atoms. These alkyl groups
 may be derived from long-chain fatty acids, such as
 hydrogenated tallow fatty acid. Tallow fatty acid gives
 rise to quaternary compounds wherein R₁ and R₂ predominantly
 have from 16 to 18 carbon atoms. Examples include ditallow
 dimethyl ammonium chloride, ditallow dimethyl ammonium
 methyl sulfate, dihexadecyl dimethyl ammonium chloride,
 di(hydrogenated tallow) dimethyl ammonium chloride,
 dioctadecyl dimethyl ammonium chloride, dieicosyl dimethyl
 ammonium chloride, didocosyl dimethyl ammonium chloride,
 di(hydrogenated tallow) dimethyl ammonium acetate,

-30-

dihexadecyl dimethyl ammonium chloride, dihexadecyl dimethyl ammonium acetate, ditallow dipropyl ammonium phosphate, ditallow dimethyl ammonium nitrate, di(coconutalkyl) dimethyl ammonium chloride, and stearyl dimethyl benzyl ammonium chloride. Preferred quaternary ammonium salts useful herein include ditallow dimethyl ammonium chloride, dicetyl dimethyl ammonium chloride, stearyl dimethyl benzyl ammonium chloride, cetyl trimethyl ammonium chloride, tricetyl methyl ammonium chloride, and mixtures thereof.

Di(hydrogenated tallow) dimethyl ammonium chloride (Quaternium-18) is a particularly preferred quaternary ammonium salt, and is available from the Sherex Chemical Company, Inc. as Adogen 442 and Adogen 442-100P.

Salts of primary, secondary and tertiary fatty amines may also be used as a cationic surfactant. The alkyl groups of such amines preferably have from 12 to 22 carbon atoms, and may be substituted or unsubstituted. Secondary and tertiary amines are preferred; and tertiary amines are particularly preferred. Examples of useful amines include stearamido propyl dimethyl amine, diethyl amino ethyl stearamine, dimethyl stearamine, dimethyl soyamine, soyamine, myristyl amine, tridecyl amine, ethyl stearylamine, N-tallowpropane diamine, ethoxylated (5 moles E.O.) stearylamine, dihydroxy ethyl stearylamine, and arachidylbehenylamine. Suitable amine salts include the

-31-

halogen, acetate, phosphate, nitrate, citrate, lactate and alkyl sulfate salts. Examples include stearylamine hydrochloride, soyamine chloride, stearylamine formate, N-tallowpropane diamine dichloride and stearamidopropyl dimethylamine citrate. Useful cationic amine surfactants are also disclosed in U.S. Pat. No. 4,275,055 (Nachtigal et al., issued Jun. 23, 1982).

Water is an essential ingredient in the conditioner formulations. Water is added as the last step in preparing the conditioner, using an amount sufficient to bring (q.s.) the mixture to 100%.

Optional ingredients in the conditioner formulations include silicone conditioning agents that may be used for their cosmetic and rheological characteristics. Silicone oils and silicone polymers are well known conditioning agents. For example, volatile silicones, organosilicone polymers in water-alcohol mixtures, and volatile silicone fluids are disclosed in U.S. Patent No. 5,292,502, supra.

The formulation may include one or more silicones disclosed for use in the shampoo formulations supra. These silicones include volatile and non-volatile polyalkyl siloxanes, polyalkylaryl siloxanes, and mixtures thereof.

-32-

They may be used at levels from about 0.2% to about 5% of the final formulation.

As with shampoos, the higher viscosity silicone gums of the siloxanes disclosed supra are preferred. These gums are rigid, as opposed to a fluid, with high molecular weights of from about 200,000 to about 1,000,000 and viscosities from about 100,000 cp to about 150,000,000 cp at 25°. Most preferred are the polydimethylsiloxane gums.

Often a significant amount of the lipid material in the conditioner is deposited on the hair, leaving it greasy. The conditioner formulations may, therefore, incorporate silicone copolyols to provide optimum conditioning benefits with the anti-lice treatment. See European Patent Application 155,806, published Sep. 25, 1985.

The silicone copolyols are polyalkylene oxide modified dimethylpolysiloxanes, herein referred to as a "dimethicone copolyols" that act as an emulsifier and reduce the deposition of the vehicle materials (lipid materials and/or cationic surfactants) on the hair. Useful dimethicone copolyols are also disclosed in U.S. Patent No. 5,292,504, supra.

-33-

The silicone copolyol is generally present at a level of from about 0.1% to about 10%, preferably from about 0.1% to about 2%, of the final formulation.

5 Dimethicone copolyols are preferred for this use. Dow Corning 190 Silicone Surfactant is a preferred dimethicone copolyol.

10 The formulations may also contain components that modify the physical and performance characteristics of the conditioning product. Such components include additional surfactants, salts, buffers, thickeners, solvents, opacifiers, pearlescent aids, preservatives, fragrance, colorants, dyes, pigments, chelators, sunscreens, vitamins, and medicinal agents. Examples of these types of components
15 are disclosed in U.S. Pat. No. 4,387,090 (Bolich, Jr., issued Jun. 7, 1983).

20 The formulations may also contain optional surfactant materials at levels such that the total level of surfactant present in the formulation (including the cationic surfactant vehicle material, described supra) is from about 0.05% to about 5%. These optional surfactant materials may be anionic, nonionic or amphoteric. Examples
25 are cetareth-20, steareth-20, sorbitan monoesters, sodium tallow alkylsulfate, and tallow betaine. Optional

-34-

surfactant materials are described in McCutcheon's *Detergents & Emulsifiers*, supra; Schwarts et al., supra; and U.S. Pat. No. 3,929,678 supra.

5 Preferred optional surfactant materials are nonionic. Such surfactants are most commonly produced by the condensation of an alkylene oxide (hydrophilic in nature) with an organic hydrophobic compound that is usually
10 aliphatic or alkyl aromatic in nature. The length of the hydrophilic or polyalkylene moiety that is condensed with any particular hydrophobic compound can be adjusted to yield a water-soluble compound having the desired degree of balance between hydrophilic and hydrophobic elements. Such nonionic surfactants include polyethylene oxide condensates
15 of alkyl phenols, condensation products of aliphatic alcohols with ethylene oxide, condensation products of ethylene oxide with a hydrophobic base formed by condensation of propylene oxide with propylene glycol, and condensation products of ethylene oxide with the product
20 resulting from the reaction of propylene oxide and ethylene diamine. Another variety of nonionic surfactant is a non-polar nonionic surfactant, typified by the amine oxide surfactants. Preferred nonionic surfactants include cetareth-20, steareth-20 and ceteth-2.

25

-35-

Salts and buffers may also be added in order to modify the product rheology. For example, salts such as potassium chloride, ammonium chloride, and sodium chloride, may be added at levels of from about 0.001% to about 1%.

5 Buffers, such as citrate or phosphate buffers, may also be added. The present formulations as finally formulated preferably have a pH of from about 3 to about 10, most preferably from about 3 to about 7.

10 Additional conditioning components may also be incorporated into the formulations. For example, proteins may be added at levels of from about 0.1% to about 10%. Cationic proteins may also serve as surfactant vehicle materials.

15 Thickening agents are preferred optional components. Such thickeners include nonionic thickening agents that are incorporated at levels of from about 0.1% to about 8%. Such agents are polymers that exhibit viscosities exceeding about 20,000 cp at low shear (about 10^{-2} sec⁻¹). Examples are
20 polyoxyethylene, guar gum, methylcellulose, methyl hydroxypropyl cellulose, polypropyl cellulose, polypropyl hydroxyethyl cellulose, hydroxyethyl cellulose, starches and starch derivatives, and mixtures thereof. Nonionic
25 thickening agents are disclosed in U.S. Pat. No. 4,387,090 (Bolich et al., issued Jun.7, 1983).

-36-

5 The thickening agents are used to bring the viscosity of the formulation from about 10,625 cp to about 14,375 cp (as measured with a Wells-Brookfield viscometer, Model RVT DV-CP-2, DV-11, Model Cone CP-52, using ½ mL at 1 rpm at 26.7°. for 1 minute).

10 The hair conditioning formulations of this invention are generally used on the hair after all shampoo has been removed by rinsing with water.

 This invention also provides a method for treating human hair to kill and facilitate removal of lice and their eggs, comprising the steps of:

- 15 (a) applying from about 10 grams to about 30 grams of a formulation of this invention to the wet hair;
- (b) working the formulation through the hair and scalp;
- 20 (c) leaving the formulation on the hair and scalp for about 6-10 minutes; and
- (d) removing the formulation from the hair by rinsing with water.

25 Anti-lice lotions comprising a spinosyn, or a physiologically acceptable derivative or salt thereof, and a

-37-

lotion carrier are another aspect of this invention. These lotions can be applied directly onto the hair in liquid form or in spray form. They are formulated to be applied to the hair for a period of time and not immediately removed by rinsing with water.

When used in a lotion formulation, the spinosyn component is generally present at a level of from about 0.1% to about 30%, preferably from about 1% to about 10%.

In addition to the spinosyn component, the lotion formulations comprise a liquid vehicle such as alcohol, water or a mixture thereof, to assist in delivery of the spinosyn component to the hair. Suitable alcohols are monohydric alcohols such as methanol, ethanol, isopropanol, or mixtures thereof. Since alcohols can have a deleterious effect upon the stability of the formulations, water alone is most preferred as the vehicle. The vehicle is added in an amount necessary to q.s. the formulation to 100%.

-38-

The lotion formulations may include optional components to provide benefits to the hair in addition to the anti-lice activity. Optional components include:

5 preservatives and antimicrobials, such as DMDM hydantoin and tetrasodium EDTA; pH balancing agents, such as sodium citrate and citric acid; emulsifiers, such as PEG-60 castor oil; and thickeners and viscosity modifiers, such as polyvinylpyrrolidone. When included, such components

10 generally are used individually at a level from about 0.01% to about 10%.

Conditioning agents may be included to facilitate the removal of dead lice and eggs from the hair and to

15 provide good wet and dry combing. The same types of conditioning agents described in the conditioning formulations supra may be used in the lotions; these include quaternary ammonium salts, fatty amines and mixtures thereof. Conditioning agents are used at levels from about

20 0.1% to about 1%, preferably from about 0.4% to about 0.6%.

Preferred conditioning agents are quaternary ammonium salts. Preferred quaternary ammonium salts include dialkyldimethylammonium chlorides, wherein the alkyl groups

25 have from 12 to 22 carbon atoms. These alkyl groups may be derived from long-chain fatty acids, such as hydrogenated

-39-

tallow fatty acid. Tallow fatty acid gives rise to quaternary compounds wherein R_1 and R_2 predominantly have from 16 to 18 carbon atoms. Examples of quaternary ammonium salts useful in the lotion formulations include

5 di(hydrogenated) tallow dimethyl ammonium chloride, dicetyl dimethyl ammonium chloride, tricetyl methyl ammonium chloride, cetyl trimethyl ammonium chloride, stearyl dimethyl benzyl ammonium chloride, and mixtures thereof. Most preferred is dicetyl dimethyl ammonium chloride.

10 Alcohol synergizers may also be added to the lotion formulations to enhance their anti-lice activity. The alcohols used in the lotion formulations are selected from phenyl C_2 - C_6 alkanols, phenyl C_2 - C_6 diols, C_2 - C_8 alkylene diols, and mixtures thereof. These synergizers may be

15 included at levels from about 0.25% to about 10%, wherein the level of phenyl alkanols, phenyl diols, and mixtures thereof, does not exceed 5% of the formulation. Preferably, the level is about 0.5% to about 5% of the formulation, most preferably from about 2% to about 4%. A preferred

20 synergizer is hexylene glycol.

The lotion formulations are applied directly to the hair. The amount of lotion used is generally from about 10 mL to about 50 mL. The lotion is worked through the hair

25 and left on the hair for about 10 minutes, preferably about

-40-

30 minutes. The hair is then cleansed, generally with a shampoo, before rinsing with water.

The following examples illustrate the formulations of this invention:

5

Example 1

A lotion formulation is prepared as follows:

	<u>Component</u>	<u>Weight (%)</u>
	Polyvinylpyrrolidone	0.50
	DMDM hydantoin	0.20
10	Tetrasodium EDTA	0.13
	Citric acid	0.05
	PEG-60 castor oil	0.50
	Hexylene glycol	4.00
	Dicetyl dimethyl ammonium chloride	0.38
15	Spinosyn A	0.50
	Water	q.s. to 100.00

20 Add the spinosyn to a tank containing a mixture of PEG-60 castor oil, hexylene glycol, propylene glycol and dicetyl dimethyl ammonium chloride at between 35° to 38°. In a second tank, mix polyvinyl pyrrolidone, DMDM hydantoin, tetrasodium EDTA and citric acid and bring the mixture to a temperature between 35° to 38°. Add the contents of the first tank to the second tank and mix until uniform. Cool
25 the mixture to about 27°, and empty into storage drums.

Example 2

-41-

A lotion formulation is prepared using the procedure described in Example 1, but with the following formula.

5	<u>Component</u>	<u>Weight (%)</u>
	Polyvinylpyrrolidone	0.50
	DMDM hydantoin	0.20
	Tetrasodium EDTA	0.13
	Citric acid	0.05
10	PEG-60 castor oil	0.50
	Hexylene glycol	2.00
	Propylene glycol	2.00
	Dicetyl dimethyl ammonium chloride	0.38
	Spinosad	0.25
15	Water	q.s. to 100.00

Example 3

20 A lotion formulation is prepared by the procedure described in Example 1, but with the following formula:

	<u>Component</u>	<u>Weight (%)</u>
25	Polyvinylpyrrolidone	0.50
	DMDM hydantoin	0.20
	Tetrasodium EDTA	0.13
	Citric acid	0.05
	Isopropanol	1.00
	PEG-60 castor oil	0.50
30	Hexylene glycol	4.00
	Dicetyl dimethyl ammonium chloride	0.60
	Spinosyn component	0.10
	Water	q.s. to 100.00

35 To control a lice infestation, the lotion formulations of Examples 1-3 are applied to the hair and

-42-

left on for at least $\frac{1}{2}$ hour before being removed by shampooing or rinsing.

Example 4

A shampoo formulation is prepared as follows:

5	<u>Component</u>	<u>Weight (%)</u>
	Ammonium laureth sulfate	10.40
	Ammonium lauryl sulfate	9.50
	Coconut monoethanolamide	4.00
	Ethylene glycol distearate	3.00
10	DMDM hydantoin	0.20
	Monosodium phosphate	0.10
	Disodium phosphate	0.25
	Citric acid	0.07
	Ammonium xylenesulfonate	1.58
15	Spinosyn A	0.50
	Water	q.s. to 100.00

Add the ammonium lauryl sulfate to a tank and heat to between about 66° to about 69°. While maintaining this temperature, add an aqueous solution of mono-sodium phosphate and then an aqueous solution of disodium phosphate. Upon reaching 69°, add the ammonium xylenesulfonate to the mixture and heat to from about 74° to 77°; add the coconut monoethanolamide, mixing until well dispersed, the ethylene glycol distearate and about 4.5% of the water. Continue mixing until homogeneous and cool mixture to about 41°. Pump the mixture into a second tank and add the ammonium laureth sulfate, DMDM hydantoin, and aqueous solution of citric acid. Add the a spinosyn to the

-43-

second tank and q.s. to 100% with water. Mix thoroughly, cool to about 27°, and pump the mixture into storage drums.

Example 5

5 A shampoo formulation is prepared as follows:

	<u>Component</u>	<u>Weight</u>
	Ammonium laureth sulfate	14.15
	Ammonium lauryl sulfate	3.14
	Coconut monoethanolamide	3.00
10	Ethylene glycol distearate	3.00
	Silicone gum ¹	0.50
	Dimethicone fluid (350 cp)	0.50
	Tricetyl methyl ammonium chloride	0.29
	Cetyl alcohol	0.42
15	Stearyl alcohol	0.18
	DMDM hydantoin	0.20
	Sodium chloride	0.90
	Ammonium chloride	0.05
	Ammonium xylenesulfonate	1.25
20	Spinosad	0.40
	Water	q.s. to 100.00

¹Silicone gum available from The General Electric Co. as SE-30 or SE-76 Gum.

25

Add approximately 0.5% of the ammonium laureth sulfate and the dimethicone to a container, and mix for approximately 30 minutes. Add approximately 2% ammonium laureth sulfate to a processing tank and heat to 68° to 71°.

30 Add about 0.12% stearyl alcohol, about 0.06% of cetyl alcohol, and the contents of the first container to the processing tank. Mix until uniform, maintaining the mixture between 68° and 71°. To a second processing tank, add

-44-

ammonium lauryl sulfate and heat to about 71°. While maintaining this temperature, add 0.05% ammonium chloride, about 18% water, ammonium xylenesulfonate and the remainder of the stearyl and cetyl alcohols. Add coconut monoethanolamide, tricetyl methyl ammonium chloride, ethylene glycol distearate, approximately half the DMDM hydantoin and the contents of the first tank to the second tank while maintaining a temperature of about 77°. Mix until homogenous and then cool to about 41°. Pump to a third tank and add the remainder of the ammonium laureth sulfate, DMDM hydantoin, and sodium chloride. Add the spinosyn to the mixture and q.s. to 100% with water. Mix thoroughly, cool to about 27°, and pump the mixture into storage drums.

Example 6

A shampoo formulation is prepared as follows:

<u>Component</u>	<u>Weight %</u>
Ammonium laureth sulfate	12.81
Ammonium lauryl sulfate	9.10
Coconut monoethanolamide	2.30
Isostearyl ethylmidonium ethosulfate	1.25
DMDM hydantoin	0.20
Monosodium pohsophate	0.50
Disodium phosphate	0.38
Sodium chloride	0.04
Citric acid	0.10

-45-

Ammonium xylenesulfonate	1.35
Spinosyn component	0.56
Water	q.s. to 100.00

5 Add about 6.5% of the water and the ammonium
laureth sulfate to a mixing tank and heat the mixture to
about 35°. While maintaining this temperature, add the
following components individually in sequence, mixing so
that each component is well mixed into the batch: ammonium
10 lauryl sulfate, ammonium xylenesulfonate, monosodium
phosphate, disodium phosphate, DMDM hydantoin, sodium
chloride, a solution of citric acid and water, a solution of
coconut diethanolamide and isostearyl ethylmidonium
ethosulfate. Add the spinosyn to the mixture, and q.s. to
15 100% with water. Mix thoroughly, cool to about 27°, and
pump the mixture into storage drums.

Example 7

20 A conditioner formulation of this invention is
prepared as follows:

<u>Component</u>	<u>Weight %</u>
Cetyl alcohol	1.00
Stearyl alcohol	0.72
DMDM hydantoin	0.20
Hydroxyethyl cellulose	0.50
Quaternium-18	0.85
Ceteareth-20	0.35
Stearalkonium chloride	0.85

-46-

	Glyceryl monostearate	0.25
	Citric acid	0.08
	Silicone gum ¹	0.30
	Cyclomethicone fluid	1.70
5	Spinosyn A	1.00
	Water	q.s. to 100.00

¹ Silicone gum available from The General Electric Co. as SE-30 or SE-76 Gum.

10 Combine all components, except the DMDM hydantoin, citric acid, silicone gum, cyclomethicone, and a spinosyn, in a processing tank and heat the mixture to about 88°. After the solution is thoroughly mixed, cool it to
15 approximately 48°. In a separate tank, premix the silicone gum and cyclomethicone, with heat and agitation to form a gum solution. Add the spinosyn to this mixture. Add the gum solution and all the remaining components, and q.s. with water. Mix thoroughly, cool to about 27°, and pump the
20 mixture into storage drums.

Example 8

25 A conditioner formulation is prepared by the procedure described in Example 7, but using the following formula:

	<u>Component</u>	<u>Weight %</u>
	Cetyl alcohol	1.00
30	Stearyl alcohol	0.72

-47-

	DMDM hydantoin	0.20
	Hydroxyethyl cellulose	0.50
	Quaternium-18	0.85
5	Ceteareth-20	0.35
	Stearamidopropyl dimethyl amine (SAPDMA)	0.50
	Glyceryl monostearate	0.25
	Citric acid	0.08
	Sodium Citrate	0.05
	Stearoxydimethicone	0.10
10	Silicone gum ¹	0.05
	Cyclomethicone fluid	1.70
	Spinosyn component	1.00
	Water	q.s. to 100.00

15 ¹ Silicone gum available from The General
Electric Co. as SE-30 or SE-76 Gum.

 This conditioner anti-lice product is made in a
manner similar to that described in Example 7.

Example 9

Efficacy of Shampoo Formulations.

 Shampoo formulations containing various
concentrations of spinosad were used in this study. The
25 formulations were prepared by wet milling for 30 minutes
enough technical grade spinosad into a commercially
available shampoo (Johnson's ® Baby Shampoo, Moisturizing
Formula with Honey and Vitamin E, Johnson & Johnson Consumer
Products, Inc.) to form a 10% stock spinosad/shampoo
30 mixture. This mixture was diluted with additional shampoo
to prepare the following spinosad concentrations: 10% (used

-48-

as originally prepared), 1%, 0.1% and 0.01% spinosad/shampoo (w/w).

5 The four concentrations of spinosad in shampoo and
a control of tap water were tested against adult human body
lice (*Pediculus humanus humanus*) according to a standard
test, ASTM Standard E 938-83 (Reapproved 1988), that is
available from the American Society for Testing and
Materials, 100 Barr Harbor Drive, West Conshohocken,
10 Pennsylvania USA [<http://www.astm.org/>]. In this test, 25 adult
lice were immersed in each of the four shampoo
concentrations for 10 minutes, then washed in water for 1
minute and rinsed in water for another minute. In the
control group, 25 adult lice (*Pediculus humanus humanus*)
15 were immersed in tap water for 10 minutes, then washed in
water for 1 minute, and rinsed in water for another minute.
A total of 5 trials were performed.

20 After one hour, the lice were examined to
determine the knockdown number. "Knockdown" was measured as
the rather quick (within a matter of one minute)
immobilization of insect activity which leads from a
moribund state to a state of kill. After 24 hours, the lice

-49-

were again examined to determine the number killed. The results of this study are summarized in Table 1.

5 **Table 1: Comparison of Pediculicidal Effects of Spinosad in Shampoo at Various Concentrations**

Mortality Data

		+1 hour	+24 hours
10	Formulation	% Knockdown	% Mortality
	Control	0.2	1.0
	10% spinosad/shampoo	96.6	100.0
	1% spinosad/shampoo	48.0	100.0
	0.1% spinosad/shampoo	19.8	97.4
15	0.01% spinosad/shampoo	14.7	35.5

20 The study showed that shampoo formulations containing 1% and 10% spinosad were highly effective pediculicides, providing a +24 hour mortality of 100%. The 10% concentration gave the quickest knockdown effect (96.6% at +1 hour), and even the 1% concentration provided a knockdown rate of 48%. The 0.1% spinosad/shampoo formulation was also an effective pediculicide, providing 25 nearly 100% mortality after 24 hours.

-50-
CLAIMS

1. A formulation for controlling a lice infestation in a human comprising as an active ingredient
5 a spinosyn, or a physiologically acceptable derivative or salt thereof, associated with a physiologically acceptable carrier.
2. A formulation as claimed in Claim 1 that
10 is a hair-care formulation.
3. A formulation as claimed in Claims 1 or 2 that is a shampoo.
4. A formulation as claimed in any one of
15 Claims 1, 2 or 3 which additionally comprises a surfactant.
5. A formulation as claimed in any one of
20 Claims 3 or 4 which comprises
- (a) from about 0.1% to about 2.5% of a spinosyn, or a physiologically acceptable derivative or salt thereof;
 - (b) from about 5% to about 30% of a
25 synthetic surfactant;
 - (c) from about 1% to about 7% of an amide; and
 - (d) water.
6. A formulation as claimed in any one of
30 Claims 1 to 5 which additionally comprises from about 1% to about 10% of a non-volatile silicone material.
7. A formulation as claimed in any one of
35 Claims 1 to 6 which additionally comprises from about 0.5% to about 5% of a suspending agent.

-51-

8. The use of a spinosyn, or a physiologically acceptable derivative or salt thereof, or a formulation containing either entity, for the manufacture of a medicament for controlling lice in a human.

9. The use of Claim 8 wherein the louse is *Pediculus humanus capitis* or *Pediculus humanus humanus*.

10. The use of Claim 10 wherein the louse is *Pthirus pubis*.

11. A method for controlling a lice infestation in a human comprising topically administering a formulation as claimed in any one of Claims 1 to 4 to the human.

12. The method of Claim 11 wherein the lice infestation is *Pediculus humanus capitis*.

13. The method of Claim 11 wherein the lice infestation is *Pediculus humanus humanus*.

14. The method of Claim 11 wherein the lice infestation is *Pthirus pubis*.

15. A method for treating human hair to kill and facilitate removal of lice and their eggs comprising the steps of:

(a) applying from about 10 g to about 30 g of a formulation comprising a spinosyn, or a physiologically acceptable derivative or salt thereof, and a physiologically acceptable carrier to wet hair;

(b) working the formulation through the hair and scalp

-52-

(c) leaving the formulation on the hair and scalp for about 6 - 10 minutes; and

(d) removing the formulation from the hair by rinsing with water.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/335	A3	(11) International Publication Number: WO 00/01347 (43) International Publication Date: 13 January 2000 (13.01.00)
(21) International Application Number: PCT/US99/13925 (22) International Filing Date: 21 June 1999 (21.06.99) (30) Priority Data: 60/091,658 2 July 1998 (02.07.98) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): SNYDER, Daniel, Earl [US/US]; 7348 Yorkshire Boulevard, Indianapolis, IN 46229 (US). (74) Agents: GRAY, Edouard, P. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AE, AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 23 March 2000 (23.03.00)
(54) Title: FORMULATIONS FOR CONTROLLING HUMAN LICE (57) Abstract Safer pediculicidal formulations comprising a spinosyn, or a physiologically acceptable derivative or salt thereof, and a physiologically acceptable carrier, and methods of controlling lice infestations in a human with these formulations are provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/13925

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/335

US CL :514/450

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/450

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS AND CAS ONLINE: spinosyn a, spinosyn d, spinosad with lice, louse, pediculus, pthirus, insect?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,571,901 A (BOECK et al) 05 November 1996, abstract, col. 1, lines 50-67, col. 30, line 8 - col. 32, line 36.	1-3, 8-10, 15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but which does not understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

01 AUGUST 1999

Date of mailing of the international search report

07 FEB 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer


KIMBERLY JORDAN

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/13925

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 4-7, 11-14
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.